

A CHARACTERISATION OF LIE ALGEBRAS AMONGST ALTERNATING ALGEBRAS

XABIER GARCÍA-MARTÍNEZ AND TIM VAN DER LINDEN

ABSTRACT. Let \mathbb{K} be an infinite field. We prove that if a variety of alternating \mathbb{K} -algebras—not necessarily associative, where $xx = 0$ is a law—is locally algebraically cartesian closed, then it must be a variety of Lie algebras over \mathbb{K} . In particular, $\text{Lie}_{\mathbb{K}}$ is the largest such. Thus, for a given variety of alternating \mathbb{K} -algebras, the Jacobi identity becomes equivalent to a categorical condition: it is a law in \mathcal{V} if and only if \mathcal{V} is a subvariety of a locally algebraically cartesian closed variety of alternating \mathbb{K} -algebras. This is based on a result saying that an algebraically coherent variety of alternating \mathbb{K} -algebras is either a variety of Lie algebras or a variety of antiassociative algebras over \mathbb{K} .

1. INTRODUCTION

The aim of this article is to prove that, if a variety of alternating algebras—not necessarily associative, where $xx = 0$ is a law—over an infinite field admits *algebraic exponents* in the sense of James Gray’s Ph.D. thesis [15], so when it is *locally algebraically cartesian closed* (or (LACC) for short, see [17, 7]), then it must necessarily be a variety of Lie algebras. Since, as shown in [16], the category $\text{Lie}_{\mathbb{K}}$ of Lie algebras over a commutative unitary ring \mathbb{K} is always (LACC), this condition may be used to characterise Lie algebras amongst alternating algebras.

The only other non-abelian “natural” examples of locally algebraically cartesian closed semi-abelian [20] categories we currently know of happen to be categories of group objects in a cartesian closed category [17], namely

- (1) the category \mathbf{Gp} of groups itself;
- (2) the category \mathbf{XMod} of crossed modules, which are the group objects in the category \mathbf{Cat} of small categories [22, 24]; and
- (3) the category $\mathbf{Hopf}_{\mathbb{K}, \text{coc}}$ of cocommutative Hopf algebras over a field \mathbb{K} of characteristic zero [14, 10], the group objects in the category $\mathbf{CoAlg}_{\mathbb{K}, \text{coc}}$ of cocommutative coalgebras over \mathbb{K} .

At first with our project we hoped to remedy this situation by finding further examples of (LACC) categories of (not necessarily associative) algebras. However, all of our attempts at constructing such new examples failed. Quite unexpectedly, in the end we managed to prove that, at least when the field \mathbb{K} is infinite, amongst those algebras which are alternating, *there are no other examples*: the condition (LACC) implies that the Jacobi identity holds. Thus, in the context of alternating algebras, the Jacobi identity is characterised in terms of a purely categorical condition. This is the subject of Section 2.

2010 *Mathematics Subject Classification.* 08C05, 17A99, 18B99, 18A22, 18D15.

Key words and phrases. Lie algebra; antiassociative, alternating algebra; algebraically coherent, locally algebraically cartesian closed, semi-abelian category; algebraic exponentiation.

This work was partially supported by Ministerio de Economía y Competitividad (Spain), grant MTM2016-79661-P. The first author was also supported by Xunta de Galicia, grant GRC2013-045 (European FEDER support included), by an FPU scholarship of the Ministerio de Educación, Cultura y Deporte (Spain) and by a Fundación Barrié scholarship. The second author is a Research Associate of the Fonds de la Recherche Scientifique–FNRS.

We do not know what happens when the algebras considered are not alternating. The category of Leibniz algebras is not (LACC), so at least one of the implications in our characterisation fails in that case. We make a few additional observations in Section 3, and hope to study this question in future work.

1.1. Cartesian closedness. *Algebraic exponentiation* is a categorical-algebraic version of the concept of *exponentiation* familiar from set theory, linear algebra, topology, etc. In its most basic form, exponentiation amounts to the task of equipping the set $\text{Hom}_{\mathcal{C}}(X, Y)$ of morphisms from X to Y with a suitable structure making it an object Y^X in the category \mathcal{C} at hand.

Depending on the given category \mathcal{C} , this may or may not be always possible. A category with binary products \mathcal{C} is said to be **cartesian closed** when for every object X the functor $X \times (-): \mathcal{C} \rightarrow \mathcal{C}$ admits a right adjoint $(-)^X: \mathcal{C} \rightarrow \mathcal{C}$, so that for all Y and Z in \mathcal{C} , the set $\text{Hom}_{\mathcal{C}}(X \times Z, Y)$ is isomorphic to $\text{Hom}_{\mathcal{C}}(Z, Y^X)$. In particular then, an object Y^X exists for all X and Y ; see [24] for further details.

The category **Set** of sets is cartesian closed, with Y^X the set $\text{Hom}_{\text{Set}}(X, Y)$ of functions from X to Y . Also the category **Cat** of small categories is cartesian closed. The category Y^X has functors $X \rightarrow Y$ as objects, and natural transformations between them as morphisms. For any commutative ring \mathbb{K} , the category $\text{CoAlg}_{\mathbb{K}, \text{coc}}$ of cocommutative coalgebras over \mathbb{K} is cartesian closed by a result in [1].

1.2. Closedness in general. The categories occurring in algebra are seldom cartesian closed. The concept of closedness has thus been extended in several different directions. One option is to replace the cartesian product by some other product, such as for instance the tensor product $\otimes_{\mathbb{K}}$ when \mathcal{C} is the category $\text{Vect}_{\mathbb{K}}$ of vector spaces over a field \mathbb{K} . In that case the result is the well-known tensor/hom adjunction, where the object Y^X in the isomorphism $\text{Hom}_{\mathbb{K}}(X \otimes_{\mathbb{K}} Z, Y) \cong \text{Hom}_{\mathbb{K}}(Z, Y^X)$ is the set of \mathbb{K} -linear maps $\text{Hom}_{\mathbb{K}}(X, Y)$ with the pointwise \mathbb{K} -vector space structure.

1.3. An alternative approach. Another option, fruitful in non-abelian algebra, is to keep the cartesianness aspect of the condition, but to make it algebraic in an entirely different way [15, 17, 7]. To do this, we first need to understand what is *local* cartesian closedness by reformulating the condition in terms of slice categories. Here we follow Section A1.5 of [21].

1.4. Local cartesian closedness. Let \mathcal{C} be any category. Given an object B of \mathcal{C} , we write $(\mathcal{C} \downarrow B)$ for the **slice category** or **category of objects over B** in which an object x is an arrow $x: X \rightarrow B$ in \mathcal{C} , and a morphism $f: x \rightarrow y$ is a commutative triangle

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \searrow x & \swarrow y \\ & B & \end{array}$$

in \mathcal{C} , so that $y \circ f = x$.

Assuming now that \mathcal{C} is finitely complete, given a morphism $a: A \rightarrow B$, we write

$$a^*: (\mathcal{C} \downarrow B) \rightarrow (\mathcal{C} \downarrow A)$$

for the **change-of-base functor** which takes an arrow $x: X \rightarrow B$ in \mathcal{C} and sends it to its pullback $a^*(x)$ as in the diagram

$$\begin{array}{ccccc} A \times_B X & \xrightarrow{\quad} & X & & \\ a^*(x) \downarrow & \lrcorner & \downarrow x & & \\ A & \xrightarrow{a} & B & & \end{array}$$

If B is the terminal object 1 of \mathcal{C} then $(\mathcal{C} \downarrow B) = (\mathcal{C} \downarrow 1) \cong \mathcal{C}$. Any object A of \mathcal{C} now induces a unique morphism $a = !_A: A \rightarrow 1$, and the functor $!_A^*: \mathcal{C} \rightarrow (\mathcal{C} \downarrow A)$ sends an object Y to the product $A \times Y$ (considered together with its projection to A). It is easily seen that the category \mathcal{C} is cartesian closed if and only if for every X in \mathcal{C} , the functor $!_X^*$ admits a right adjoint.

A category with finite limits \mathcal{C} is said to be **locally cartesian closed** or **(LCC)** when for *every* morphism $a: A \rightarrow B$ in \mathcal{C} the change-of-base functor a^* has a right adjoint. Equivalently, all slice categories $(\mathcal{C} \downarrow B)$ are cartesian closed—so that \mathcal{C} is cartesian closed, *locally over* B , for all B in \mathcal{C} . This condition is stronger than cartesian closedness (the case $B = 1$); examples include any Grothendieck topos, in particular the category of sets, while for instance [12] the category **Cat** is not (LCC), even though it is cartesian closed.

1.5. Categories of points. We may now modify the concept of (local) cartesian closedness in such a way that it applies to algebraic categories. The idea is that, where *slice categories* are useful in non-algebraic settings, in algebraic categories a similar role may be played by *categories of points*.

Let \mathcal{C} be any category. Given an object B of \mathcal{C} , we write $\text{Pt}_B(\mathcal{C})$ for the **category of points over** B in which an object (x, s) is a split epimorphism $x: X \rightarrow B$ in \mathcal{C} , together with a chosen section $s: B \rightarrow X$, so that $x \circ s = 1_B$. Given two points $(x: X \rightarrow B, s: B \rightarrow X)$ and $(y: Y \rightarrow B, t: B \rightarrow Y)$ over B , a morphism between them is an arrow $f: X \rightarrow Y$ in \mathcal{C} satisfying $y \circ f = x$ and $f \circ s = t$.

Change of base is done as for slice categories: since sections are preserved, given any morphism $a: A \rightarrow B$ in a finitely complete category \mathcal{C} , we obtain a functor

$$a^*: \text{Pt}_B(\mathcal{C}) \rightarrow \text{Pt}_A(\mathcal{C}).$$

1.6. Protomodular and semi-abelian categories. A finitely complete category \mathcal{C} is said to be **Bourn protomodular** [3, 5, 2] when each of the change-of-base functors $a^*: \text{Pt}_B(\mathcal{C}) \rightarrow \text{Pt}_A(\mathcal{C})$ reflect isomorphisms. If \mathcal{C} is a pointed category, then this condition may be reduced to the special case where A is the zero object and $a = !_B: 0 \rightarrow B$ is the unique morphism. The pullback functor $!_B^*: \text{Pt}_B(\mathcal{C}) \rightarrow \mathcal{C}$ then sends a split epimorphism to its kernel. Hence, protomodularity means that the **Split Short Five Lemma** holds: suppose that in the commutative diagram

$$\begin{array}{ccccc} K & \xrightarrow{k} & X & \begin{smallmatrix} \xrightarrow{x} \\ \xleftarrow{s} \end{smallmatrix} & B \\ g \downarrow & & f \downarrow & & \parallel \\ L & \xrightarrow{l} & Y & \begin{smallmatrix} \xrightarrow{y} \\ \xleftarrow{t} \end{smallmatrix} & B, \end{array}$$

the morphism k is the kernel of x and l is the kernel of y , while f is a morphism of points $(x, s) \rightarrow (y, t)$; if now g is an isomorphism, then f is also an isomorphism.

A pointed protomodular category which is Barr exact and has finite coproducts is called a **semi-abelian** category [20]. This concept unifies earlier attempts (including, for instance, [19, 13, 28]) at providing a categorical framework that extends the context of abelian categories to encompass non-additive categories of algebraic structures such as groups, Lie algebras, loops, rings, etc. In this setting, the basic lemmas of homological algebra—the 3×3 Lemma, the *Short Five Lemma*, the *Snake Lemma*—hold [5, 2], and may be used to study, say, (co)homology, radical theory, or commutator theory for those non-additive structures.

In a semi-abelian category, any point (x, s) with its induced kernel k as above gives rise to a split extension, since x is also the cokernel of k , so that (k, x) is a short exact sequence. By the results in [8], split extensions are equivalent to so-called *internal actions* by means of a semi-direct product construction. Through this

equivalence, there is a unique internal action $\xi: B\flat K \rightarrow K$ such that $X \cong K \rtimes_{\xi} B$. Without going into further details, let us just mention here that the object $B\flat K$ is the kernel of the morphism $(1_B \ 0): B + K \rightarrow B$, that the functor $B\flat(-): \mathcal{C} \rightarrow \mathcal{C}$ is part of a monad, and that an internal B -action is an algebra for this monad. The category $\mathbf{Pt}_B(\mathcal{C})$ is monadic over \mathcal{C} , and its equivalence with the category of $B\flat(-)$ -algebras bears witness of this fact.

1.7. Examples. All *Higgins varieties of Ω -groups* [18] are semi-abelian, which means that any pointed variety of universal algebras whose theory contains a group operation is an example. In particular, we find categories of all kinds of (not necessarily associative) algebras over a ring as examples, next to the categories of groups, crossed modules, and groups of a certain nilpotency or solvability class. Other examples include the categories of Heyting semilattices, loops, compact Hausdorff groups and the dual of the category of pointed sets [20, 2].

1.8. Algebraic cartesian closedness and the condition (LACC). A category with finite limits \mathcal{C} is said to be **locally algebraically cartesian closed** or **(LACC)** when for every morphism $a: A \rightarrow B$ in \mathcal{C} , the change-of-base functor $a^*: \mathbf{Pt}_B(\mathcal{C}) \rightarrow \mathbf{Pt}_A(\mathcal{C})$ has a right adjoint [15]. This condition is much stronger than **algebraic cartesian closedness** or **(ACC)** which is the case $B = 1$.

When a semi-abelian category is (locally) algebraically cartesian closed, this has some interesting consequences [17, 7, 10]. For one thing, (ACC) is equivalent to the condition that every monomorphism in \mathcal{C} admits a centraliser. The property (LACC) implies categorical-algebraic conditions such as *peri-abelianness* [6], *strong protomodularity* [4], the *Smith is Huq* condition [25], *normality of Higgins commutators* [11], and *algebraic coherence*. We come back to the latter condition (which implies all the others mentioned) in detail, in Subsection 1.10 below.

The condition (ACC) is relatively weak, and has all *Orzech categories of interest* [28] for examples. In comparison, (LACC) is very strong: as mentioned above, we have groups, Lie algebras, crossed modules, and cocommutative Hopf algebras over a field of characteristic zero as “natural” semi-abelian examples, next to all abelian categories. An example of a slightly different kind—because it is non-pointed—is any category of groupoids with a fixed object of objects [7].

In what follows, we shall need the following characterisation of (LACC), valid in semi-abelian varieties of universal algebras. Instead of checking that all change-of-base functors $a^*: \mathbf{Pt}_B(\mathcal{V}) \rightarrow \mathbf{Pt}_A(\mathcal{V})$ have a right adjoint, it suffices to check that *some* change-of-base functors preserve binary sums.

Theorem 1.9. *For a semi-abelian variety of universal algebras \mathcal{V} , the following are equivalent:*

- (i) \mathcal{V} is locally algebraically cartesian closed;
- (ii) for all B in \mathcal{V} , the pullback functor $i_B^*: \mathbf{Pt}_B(\mathcal{V}) \rightarrow \mathcal{V}$ preserves all colimits;
- (iii) for all B in \mathcal{V} , the functor i_B^* preserves binary sums;
- (iv) the canonical comparison $(B\flat_{i_X} \ B\flat_{i_Y}): B\flat X + B\flat Y \rightarrow B\flat(X + Y)$ is an isomorphism for all B, X and Y in \mathcal{V} .

Proof. This combines Theorem 2.9, Theorem 5.1 and Proposition 6.1 in [17]. \square

Via the equivalence between split extensions and internal actions, condition (ii) means that the forgetful functor from the category of B -actions in \mathcal{V} to \mathcal{V} preserves all colimits. Hence in this varietal context, (LACC) amounts to the property that colimits in the category of internal B -actions in \mathcal{V} are independent of the acting object B , and computed in the base category \mathcal{V} .

1.10. Algebraic coherence. The concept of an **algebraically coherent** category was introduced in [10] with the aim in mind of having a condition with strong categorical-algebraic consequences such as the ones mentioned above for (LACC), while at the same time keeping all *Orzech categories of interest* as examples. It is to *coherence* in the sense of topos theory [21, Section A1.4] what *algebraic cartesian closedness* is to *cartesian closedness*: a condition involving slice categories has been replaced by a condition in terms of categories of points.

The formal definition is that all change-of-base functors $a^*: \mathbf{Pt}_B(\mathcal{C}) \rightarrow \mathbf{Pt}_A(\mathcal{C})$ preserve jointly strongly epimorphic pairs of arrows. This is clearly weaker than asking that the a^* preserve all colimits. We shall only need the following characterisation, which is essentially Theorem 3.18 in [10]: algebraic coherence is equivalent to the condition that for all B, X and Y , the canonical comparison

$$(Bb_{\iota_X} Bb_{\iota_Y}): BbX + BbY \rightarrow Bb(X + Y)$$

from Theorem 1.9 is a regular epimorphism.

Algebraic coherence has somewhat better stability properties than (LACC). For instance, any subvariety of a semi-abelian algebraically coherent variety is still algebraically coherent. We shall come back to this in the next section.

Some examples of semi-abelian varieties which are *not* algebraically coherent are the varieties of loops, Heyting semilattices, and non-associative algebras (the category $\mathbf{Alg}_{\mathbb{K}}$ defined below).

2. MAIN RESULT

The aim of this section is to prove Theorem 2.17, which says that any (LACC) variety of alternating algebras over an infinite field \mathbb{K} is a category of Lie algebras over \mathbb{K} . On the way we fully characterise algebraically coherent varieties of alternating algebras (Theorem 2.14). This is an application of a more general result telling us that a variety of \mathbb{K} -algebras is algebraically coherent if and only if it is an *Orzech category of interest* (Theorem 2.12).

2.1. Categories of algebras and their subvarieties. Let \mathbb{K} be a field. A **(non-associative) algebra A over \mathbb{K}** is a \mathbb{K} -vector space equipped with a bilinear operation $[-, -]: A \times A \rightarrow A$, so a linear map $A \otimes A \rightarrow A$. We use the notations $[x, y] = x \cdot y = xy$ depending on the situation at hand, always keeping in mind that the multiplication need not be associative. We write $\mathbf{Alg}_{\mathbb{K}}$ for the category of algebras over \mathbb{K} with product-preserving linear maps between them. It is a semi-abelian category which is not algebraically coherent. A **subvariety** of $\mathbf{Alg}_{\mathbb{K}}$ is any equationally defined class of algebras, considered as a full subcategory \mathcal{V} of $\mathbf{Alg}_{\mathbb{K}}$.

The category of **associative algebras** over \mathbb{K} is the subvariety of $\mathbf{Alg}_{\mathbb{K}}$ satisfying $x(yz) = (xy)z$.

The category $\mathbf{Alt}_{\mathbb{K}}$ of **alternating algebras** over \mathbb{K} is the subvariety of $\mathbf{Alg}_{\mathbb{K}}$ satisfying $xx = 0$. It is easily seen that those algebras are **anticommutative**, which means that $xy = -yx$ holds. If the characteristic of the field \mathbb{K} is different from 2, then the two classes coincide.

The category $\mathbf{AAAlg}_{\mathbb{K}}$ of **antiassociative algebras** over \mathbb{K} is the subvariety of $\mathbf{Alg}_{\mathbb{K}}$ satisfying $x(yz) = -(xy)z$. We write $\mathbf{AAAAAlg}_{\mathbb{K}}$ for the category of alternating antiassociative algebras over \mathbb{K} .

The category $\mathbf{Lie}_{\mathbb{K}}$ of **Lie algebras** over \mathbb{K} is the subvariety of alternating algebras satisfying the **Jacobi identity** $x(yz) + z(xy) + y(zx) = 0$.

An algebra is **abelian** when it satisfies $xy = 0$. The subvariety of $\mathbf{Alg}_{\mathbb{K}}$ determined by the abelian algebras is isomorphic to the category $\mathbf{Vect}_{\mathbb{K}}$ of vector spaces

over \mathbb{K} . An algebra A is abelian if and only if $+: A \times A \rightarrow A$ is an algebra morphism, which makes $(A, +, 0)$ an internal abelian group, so an **abelian object** in the sense of [2].

2.2. Algebras over infinite fields. We assume that the field \mathbb{K} is infinite, so that we can use the following result (Theorem 2.3, which is Corollary 2 on page 8 of [29]). We first fix some terminology. For a given set S , a **polynomial** with variables in S is an element of the free \mathbb{K} -algebra on S . Recall that the left adjoint $\mathbf{Set} \rightarrow \mathbf{Alg}_{\mathbb{K}}$ factors as a composite of the *free magma* functor $M: \mathbf{Set} \rightarrow \mathbf{Mag}$ with the *magma algebra* functor $\mathbb{K}[-]: \mathbf{Mag} \rightarrow \mathbf{Alg}_{\mathbb{K}}$. The elements of $M(S)$ are non-associative words in the alphabet S , and the elements of $\mathbb{K}[M(S)]$, the polynomials, are \mathbb{K} -linear combinations of such words. A **monomial** in $\mathbb{K}[M(S)]$ is any scalar multiple of an element of $M(S)$. The **type** of a monomial $\varphi(x_1, \dots, x_n)$ is the element $(k_1, \dots, k_n) \in \mathbb{N}^n$ where k_i is the degree of x_i in $\varphi(x_1, \dots, x_n)$. A polynomial is **homogeneous** if its monomials are all of the same type. Any polynomial may thus be written as a sum of homogeneous polynomials, which are called its **homogeneous components**.

Theorem 2.3. [29] *If \mathcal{V} is a variety of algebras over an infinite field, then all of its laws are of the form $\phi(x_1, \dots, x_n) = 0$, where $\phi(x_1, \dots, x_n)$ is a (non-associative) polynomial, each of whose homogeneous components $\psi(x_{i_1}, \dots, x_{i_m})$ again gives rise to a law $\psi(x_{i_1}, \dots, x_{i_m}) = 0$ in \mathcal{V} .* \square

2.4. Description of $B \flat X$ in $\mathbf{Alg}_{\mathbb{K}}$. Let B and X be free \mathbb{K} -algebras. Then the object $B \flat X$, being the kernel of the morphism $(1_B \ 0): B + X \rightarrow B$, consists of those polynomials with variables in B and in X which can be written in a form where all of their monomials contain variables in X . For instance, given $b, b' \in B$ and $x \in X$, the expression $(b(xx))b'$ is allowed, but bb' is not.

2.5. The reflection to a subvariety \mathcal{V} of $\mathbf{Alg}_{\mathbb{K}}$. Let B and X be free \mathbb{K} -algebras. We write \overline{B} and \overline{X} for their respective reflections into \mathcal{V} , which are free \mathcal{V} -algebras. These induce short exact sequences in $\mathbf{Alg}_{\mathbb{K}}$ such as

$$0 \longrightarrow [X] \longrightarrow X \xrightarrow{\eta_X} \overline{X} \longrightarrow 0$$

where η_X is the unit at X of the reflection from $\mathbf{Alg}_{\mathbb{K}}$ to \mathcal{V} . We never write the right adjoint inclusion, but note that it preserves all limits. The kernel $[X]$ is a kind of relative commutator; all of its elements are laws of \mathcal{V} . Reflecting sums now, then taking kernels to the left, we obtain horizontal split exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & (B \flat X) \cap [B + X] & \longrightarrow & [B + X] & \xrightleftharpoons{\quad} & [B] \longrightarrow 0 \\ & & \downarrow \text{dotted} & & \downarrow & & \downarrow \\ 0 & \longrightarrow & B \flat X & \longrightarrow & B + X & \xrightleftharpoons{(1_B \ 0)} & B \longrightarrow 0 \\ & & \downarrow \rho_{B,X} & & \downarrow \eta_{B+X} & & \downarrow \eta_B \\ 0 & \longrightarrow & \overline{B} \flat_{\mathcal{V}} \overline{X} & \longrightarrow & \overline{B} +_{\mathcal{V}} \overline{X} & \xrightleftharpoons{(1_{\overline{B}} \ 0)} & \overline{B} \longrightarrow 0 \end{array}$$

in $\mathbf{Alg}_{\mathbb{K}}$, where the sum $\overline{B} +_{\mathcal{V}} \overline{X} \cong \overline{B + X}$ is taken in \mathcal{V} . Using, for instance, the 3×3 Lemma, it is not difficult to see that the induced dotted arrow $\rho_{B,X}$ is a surjective algebra homomorphism. In fact, the upper left square is a pullback, and we have three vertical short exact sequences. The one on the left allows us to view the elements of $\overline{B} \flat_{\mathcal{V}} \overline{X}$ as polynomials in $B \flat X$, modulo those laws which hold in \mathcal{V} that are expressible in $B \flat X$. These laws are precisely the elements of the top left intersection. We freely use this interpretation in what follows, abusing terminology and notation by making no distinction between the equivalence class of polynomials

that is an element in the quotient $\overline{B\mathfrak{b}}_{\mathcal{V}}\overline{X}$, and an element in $B\mathfrak{b}X$ which represents it.

2.6. Subvarieties of $\text{Lie}_{\mathbb{K}}$ need not be (LACC). Subvarieties of locally algebraically cartesian closed categories need no longer be such: we may take the variety of Lie algebras that satisfy $x(yz) = 0$ as an example.

Proposition 2.7. *Let \mathcal{V} be a variety of non-associative algebras in which $x(yz) = 0$ is a law. If \mathcal{V} is locally algebraically cartesian closed, then it is abelian.*

Proof. Let B , X and Y be free algebras in \mathcal{V} , respectively generated by their elements b , x and y . Then the split epimorphisms

$$((1_B \ 0): B + X \rightarrow B, \quad \iota_B: B \rightarrow B + X)$$

and

$$((1_B \ 0): B + Y \rightarrow B, \quad \iota_B: B \rightarrow B + Y)$$

correspond to the free B -actions respectively generated by x and y . Their sum in $\text{Pt}_B(\mathcal{V})$ is

$$((1_B \ 0 \ 0): B + X + Y \rightarrow B, \quad \iota_B: B \rightarrow B + X + Y).$$

Applying the kernel functor, (LACC) tells us that the canonical morphism

$$(B\mathfrak{b}_{\iota_X} \ B\mathfrak{b}_{\iota_Y}): B\mathfrak{b}X + B\mathfrak{b}Y \rightarrow B\mathfrak{b}(X + Y)$$

is an isomorphism (Theorem 1.9). When considering the sum $B\mathfrak{b}X + B\mathfrak{b}Y$ as a subobject of the coproduct $B + X + B + Y$, we write b_1 and b_2 for the generators of the two distinct copies of B ; then $(B\mathfrak{b}_{\iota_X} \ B\mathfrak{b}_{\iota_Y})$ maps the b_i to b , sends x to x and y to y .

Now $x \in B\mathfrak{b}X$ and $yb_2 \in B\mathfrak{b}Y$ are such that $x \cdot yb_2$ is sent to zero by the above isomorphism, since the law $x(yz) = 0$ holds in \mathcal{V} . As a consequence, $x \cdot yb_2$ is zero in the sum $B\mathfrak{b}X + B\mathfrak{b}Y$. Recall that $b_2 \notin B\mathfrak{b}Y$, so that yb_2 cannot be decomposed as a product of y and b_2 in $B\mathfrak{b}Y$. By Theorem 2.3, yb_2 can also not be written as a product in which more than one y or b_2 appears, unless yb_2 is zero in $B\mathfrak{b}Y$. As a consequence, $x \cdot yb_2$ can only be zero if either yb_2 is zero in $B\mathfrak{b}Y$, or $xz = 0$ is a law in \mathcal{V} . In the former case, yb_2 is zero in the sum $B + Y$, which is a free algebra on $\{b_2, y\}$; then $yz = 0$ is a law in \mathcal{V} . In either case, \mathcal{V} is abelian. \square

2.8. (Anti)associative algebras. Essentially the same argument gives us two further examples, which we shall need later on:

Proposition 2.9. *If a variety of either associative or antiassociative algebras is locally algebraically cartesian closed, then it is abelian.*

Proof. In the antiassociative case we have $x, -xb_1 \in B\mathfrak{b}X$ and $b_2y, y \in B\mathfrak{b}Y$ such that $x \cdot b_2y$ and $-xb_1 \cdot y$ are sent to the same element in $B\mathfrak{b}(X + Y)$ by the above isomorphism $(B\mathfrak{b}_{\iota_X} \ B\mathfrak{b}_{\iota_Y}): B\mathfrak{b}X + B\mathfrak{b}Y \rightarrow B\mathfrak{b}(X + Y)$.

Similarly, in the associative case, we see that $xb_1 \cdot y$ and $x \cdot b_2y$ are two distinct elements of the sum $B\mathfrak{b}X + B\mathfrak{b}Y$ which the morphism $(B\mathfrak{b}_{\iota_X} \ B\mathfrak{b}_{\iota_Y})$ sends to one and the same element of $B\mathfrak{b}(X + Y)$. \square

Lemma 2.10. *Any variety of \mathbb{K} -algebras that satisfies the law $x(xy) = 0$ is a subvariety of $\text{AAAlg}_{\mathbb{K}}$.*

Proof. Taking $x = a + b$ and $y = c$ gives us

$$0 = (a + b)((a + b)c) = a(ac) + b(ac) + a(bc) + b(bc) = b(ac) + a(bc)$$

so that $a(bc) = -b(ac)$. It follows that $(uv)w = -w(uv) = u(vw) = -u(vw)$ is a law in \mathcal{V} , and \mathcal{V} is a variety of antiassociative algebras. \square

2.11. Algebraic coherence. Theorem 2.3 gives us a characterisation of algebraic coherence for varieties of \mathbb{K} -algebras.

Theorem 2.12. *Let \mathbb{K} be an infinite field. If \mathcal{V} is a variety of non-associative \mathbb{K} -algebras, then the following are equivalent:*

- (i) \mathcal{V} is algebraically coherent;
- (ii) there exist $\lambda_1, \dots, \lambda_8$ in \mathbb{K} such that

$$\begin{aligned} z(xy) = & \lambda_1 y(zx) + \lambda_2 x(yz) + \lambda_3 y(xz) + \lambda_4 x(zy) \\ & + \lambda_5 (zx)y + \lambda_6 (yz)x + \lambda_7 (xz)y + \lambda_8 (zy)x \end{aligned}$$

is a law in \mathcal{V} ;

- (iii) \mathcal{V} is an Orzech category of interest [28].

Proof. From the results of [10] we already know that (iii) implies (i). It follows immediately from the definition of an *Orzech category of interest* that (ii) implies (iii). To see that (i) implies (ii), we take free B -actions as in the first part of the proof of Proposition 2.7 and obtain the regular epimorphism

$$(B\flat_X \ B\flat_Y): B\flat X + B\flat Y \rightarrow B\flat(X + Y).$$

Any element $b(xy)$ of $B\flat(X + Y)$ is the image through this morphism of some polynomial $\psi(b_1, x, b_2, y)$ in $B\flat X + B\flat Y$. Note that this polynomial cannot contain any monomials obtained as a product of a b_i with xy or yx . This allows us to write, in the sum $B + X + Y$, the element $b(xy)$ as

$$\begin{aligned} & \lambda_1 y(bx) + \lambda_2 x(yb) + \lambda_3 y(xb) + \lambda_4 x(by) + \lambda_5 (bx)y + \lambda_6 (yb)x + \lambda_7 (xb)y + \lambda_8 (by)x \\ & + \nu \phi(b, x, y) \end{aligned}$$

for some $\lambda_1, \dots, \lambda_8, \nu \in \mathbb{K}$, where $\phi(b, x, y)$ is the part of the polynomial in b, x and y which is not in the homogeneous component of $b(xy)$. Since $B + X + Y$ is the free \mathcal{V} -algebra on three generators b, x and y , from Theorem 2.3 we deduce that the equation in (ii) is again a law in \mathcal{V} . \square

Remark 2.13. This result may be used to prove the claim made in [10] that the category of **Jordan algebras**—commutative and such that $(xy)(xx) = x(y(xx))$ —is not algebraically coherent. Indeed, as explained in [27], it is not a *category of interest*.

In the case of alternating algebras, this characterisation becomes more precise:

Theorem 2.14. *Let \mathbb{K} be an infinite field. If \mathcal{V} is a subvariety of $\text{Alt}_{\mathbb{K}}$, then the following are equivalent:*

- (i) \mathcal{V} is algebraically coherent;
- (ii) \mathcal{V} is a subvariety of either $\text{AAAAlg}_{\mathbb{K}}$ or $\text{Lie}_{\mathbb{K}}$.

Proof. (ii) implies (i) since $\text{AAAAlg}_{\mathbb{K}}$ and $\text{Lie}_{\mathbb{K}}$ are *Orzech categories of interest* [28], so their subvarieties are algebraically coherent. To see that (i) implies (ii), we first use anticommutativity to simplify the law given in Theorem 2.12 to

$$z(xy) = \lambda y(zx) + \mu x(yz)$$

for some λ and μ in \mathbb{K} . Choosing, in turn, $y = z$ and $x = z$, we see that

- (1) either $\lambda = -1$ or $z \cdot zx = 0$ is a law in \mathcal{V} , and
- (2) either $\mu = -1$ or $x \cdot xy = 0$ is a law in \mathcal{V} .

In any of the latter cases, \mathcal{V} is a variety of antiassociative algebras by Lemma 2.10, which makes it abelian (Proposition 2.9). We are left with the situation when $\lambda = \mu = -1$, which means that the Jacobi identity is a law in \mathcal{V} , so that \mathcal{V} is a variety of Lie algebras. \square

Example 2.15. The variety of *alternating associative algebras* is an example. We have that $0 = x(yy) = (xy)y$ is a law, so that by Lemma 2.10 those algebras are antiassociative as well. It follows that $xyz = 0$ is a law. We regain a variety as in Proposition 2.7, so since it is not abelian, it cannot be (LACC).

2.16. A characterisation of Lie algebras amongst alternating algebras. The condition (LACC) eliminates one of the two options in Theorem 2.14.

Theorem 2.17. *Let \mathbb{K} be an infinite field. If \mathcal{V} is a locally algebraically cartesian closed variety of alternating \mathbb{K} -algebras, then it is a subvariety of $\text{Lie}_{\mathbb{K}}$. In other words, $\text{Lie}_{\mathbb{K}}$ is the largest (LACC) variety of alternating \mathbb{K} -algebras. Thus for any variety \mathcal{V} of alternating \mathbb{K} -algebras, the following are equivalent:*

- (i) \mathcal{V} is a subvariety of a (LACC) variety of alternating \mathbb{K} -algebras;
- (ii) the Jacobi identity is a law in \mathcal{V} .

Proof. This combines Theorem 2.14 with Proposition 2.9. □

Remark 2.18. By Proposition 2.7, the condition

- (iii) \mathcal{V} is (LACC)

is strictly stronger than the equivalent conditions (i) and (ii).

Remark 2.19. We do not know any non-abelian examples of (LACC) subvarieties of $\text{Lie}_{\mathbb{K}}$, or whether such subvarieties even exist.

3. NON-ALTERNATING ALGEBRAS

An important question which we have to leave open for now, is what happens when the algebras we consider are not alternating. We end this note with some of our preliminary findings.

3.1. Some of the results and techniques used in the previous section are valid for non-alternating algebras of course. For instance, Proposition 2.7, Proposition 2.9 and Theorem 2.12 are.

3.2. Proposition 2.7 contradicts—in spirit only, since we are working over a field—Proposition 6.9 in [17], which claims that the category of all commutative non-unitary rings satisfying $xyz = 0$ is locally algebraically cartesian closed. It turns out that the argument given in Proposition 2.7 is still valid in this case, and shows that the variety under consideration, should it be (LACC), would be abelian—which is false.

We noticed that the functor R constructed in the proof of [17, Proposition 6.9] is not well defined on morphisms. Let us give a concrete example showing this in detail. We follow the notations from [17, Proposition 6.9]. Let $B = \langle b \rangle = \mathbb{Z}[b]$ act on the commutative ring

$$X = \langle x, y \mid xx = xy = yy = 0 \rangle = \mathbb{Z}[x, y] / (xx, xy, yy)$$

by $bx = y$ and $by = 0$. Consider

$$M = \langle m, p, q \mid mp = q, mq = pq = mm = pp = qq = 0 \rangle$$

with the trivial B -action, and let $g: X \rightarrow M$ be the ring homomorphism sending x and y to m . Let $f \in R(X)$ be defined by $f(n, b) = nx + bx$. Then

$$R(g)(f)(0, b) \cdot p = (g \circ f)(0, b) \cdot p = g(y) \cdot n = mp = q \neq 0,$$

which shows that $R(g)(f)$ is not an element of $R(M)$.

3.3. The category $\text{Leib}_{\mathbb{K}}$ of **(right) Leibniz algebras** [23] over \mathbb{K} is the subvariety of $\text{Alg}_{\mathbb{K}}$ satisfying the **(right) Leibniz identity** $(xy)z = x(yz) + (xz)y$. This law is clearly equivalent to the Jacobi identity when the algebras are alternating, so that a Lie algebra is the same thing as an alternating Leibniz algebra. However, examples of non-alternating Leibniz algebras exist. Analogously, we can consider the category of **(left) Leibniz algebras**, with corresponding identity $x(yz) = (xy)z + y(xz)$. Both categories are of course equivalent.

We do not know whether Theorem 2.17 extends to the non-alternating case. What is certain, though, is that the category of Leibniz algebras is not locally algebraically cartesian closed. Indeed, using the notations of Proposition 2.7, the Leibniz identity allows us to deduce

$$\begin{cases} (xy)b = x(yb) + (xb)y \\ (xy)b = -x(by) + (xb)y \end{cases}$$

so that $x \cdot yb = -x \cdot by$ in $B + X + Y$. This means that $x \cdot yb_2$ and $-x \cdot b_2y$ are two distinct elements of $BbX + BbY$ which are sent to the same element of $Bb(X + Y)$ by the morphism $(Bb\iota_X \ Bb\iota_Y)$. Hence this morphism cannot be an isomorphism, and $\text{Leib}_{\mathbb{K}}$ is not (LACC).

We may ask ourselves what happens in the “intersection” between right and left Leibniz algebras. They are called **symmetric Leibniz algebras** and, as shown in [26], the chain of inclusions $\text{Lie}_{\mathbb{K}} \subseteq \text{SLeib}_{\mathbb{K}} \subseteq \text{Leib}_{\mathbb{K}}$ is strict. Doing a rearrangement of terms as in

$$b(xy) = (bx)y + x(by) = b(xy) + (by)x + x(by),$$

we see that $(by)x + x(by) = 0$. From this we may conclude that, in order to be (LACC), a variety of symmetric Leibniz algebras must either be alternating or abelian. We thus regain the known cases of Lie algebras and vector spaces.

3.4. A variety of algebras is alternating precisely when the free algebra on a single generator is abelian: $xx = 0$ is a law in the variety, if and only if the bracket vanishes on the algebra freely generated by $\{x\}$. This corresponds to the condition that *the free algebra on a single generator admits an internal abelian group structure*. This condition makes sense in arbitrary semi-abelian varieties, and we may ask ourselves whether perhaps it is implied by (LACC), as in the case of symmetric Leibniz algebras. This would allow us to drop the condition that \mathcal{V} is alternating in Theorem 2.17.

The example of crossed modules proves that this is false. In [9] it is shown that on the one hand, a crossed module $\partial: T \rightarrow G$ with action ξ admits an internal abelian group structure if and only if the groups T and G are abelian and the action ξ is trivial. On the other hand, the free crossed module on a single generator is the inclusion $\kappa: \mathbb{Z}b\mathbb{Z} \rightarrow \mathbb{Z} + \mathbb{Z}$, equipped with the conjugation action. We see that in this case the free object on one generator is not abelian, even though \mathbf{XMod} is a locally algebraically cartesian closed semi-abelian variety. However, it is not a variety of non-associative algebras of course.

3.5. Perhaps this is not the right conceptualisation, and we must think of other ways of making the law $xx = 0$ categorical. The question then becomes whether (LACC), or any other appropriate categorical-algebraic condition, would imply this new characterisation.

ACKNOWLEDGEMENTS

Thanks to James R. A. Gray, George Janelidze and Zurab Janelidze for fruitful discussions and important comments on our work. We would also like to thank

the University of Cape Town and Stellenbosch University for their kind hospitality during our stay in South Africa.

REFERENCES

- [1] M. Barr, *Coalgebras over a commutative ring*, J. Algebra **32** (1974), 600–610.
- [2] F. Borceux and D. Bourn, *Mal'cev, protomodular, homological and semi-abelian categories*, Math. Appl., vol. 566, Kluwer Acad. Publ., 2004.
- [3] D. Bourn, *Normalization equivalence, kernel equivalence and affine categories*, Category Theory, Proceedings Como 1990 (A. Carboni, M. C. Pedicchio, and G. Rosolini, eds.), Lecture Notes in Math., vol. 1488, Springer, 1991, pp. 43–62.
- [4] D. Bourn, *Normal functors and strong protomodularity*, Theory Appl. Categ. **7** (2000), no. 9, 206–218.
- [5] D. Bourn, *3×3 Lemma and protomodularity*, J. Algebra **236** (2001), 778–795.
- [6] D. Bourn, *The cohomological comparison arising from the associated abelian object*, preprint [arXiv:1001.0905](https://arxiv.org/abs/1001.0905), 2010.
- [7] D. Bourn and J. R. A. Gray, *Aspects of algebraic exponentiation*, Bull. Belg. Math. Soc. Simon Stevin **19** (2012), 823–846.
- [8] D. Bourn and G. Janelidze, *Protomodularity, descent, and semidirect products*, Theory Appl. Categ. **4** (1998), no. 2, 37–46.
- [9] P. Carrasco, A. M. Cegarra, and A. R.-Grandjeán, *(Co)Homology of crossed modules*, J. Pure Appl. Algebra **168** (2002), no. 2-3, 147–176.
- [10] A. S. Cigoli, J. R. A. Gray, and T. Van der Linden, *Algebraically coherent categories*, Theory Appl. Categ. **30** (2015), no. 54, 1864–1905.
- [11] A. S. Cigoli, J. R. A. Gray, and T. Van der Linden, *On the normality of Higgins commutators*, J. Pure Appl. Algebra **219** (2015), 897–912.
- [12] F. Conduché, *Au sujet de l'existence d'adjoints à droite aux foncteurs "image réciproque" dans la catégorie des catégories*, C. R. Acad. Sci. Paris Sér. A-B **275** (1972), A891–A894.
- [13] M. Gerstenhaber, *A categorical setting for the Baer extension theory*, Applications of Categorical Algebra, New York 1968, Proc. Sympos. Pure Math., vol. XVII, Amer. Math. Soc., Providence, R.I., 1970, pp. 50–64.
- [14] M. Gran, G. Kadjo, and J. Vercruysse, *A torsion theory in the category of cocommutative Hopf algebras*, Appl. Categ. Structures **24** (2016), no. 3, 269–282.
- [15] J. R. A. Gray, *Algebraic exponentiation in general categories*, Ph.D. thesis, University of Cape Town, 2010.
- [16] J. R. A. Gray, *Algebraic exponentiation for categories of Lie algebras*, J. Pure Appl. Algebra **216** (2012), 1964–1967.
- [17] J. R. A. Gray, *Algebraic exponentiation in general categories*, Appl. Categ. Structures **20** (2012), 543–567.
- [18] P. J. Higgins, *Groups with multiple operators*, Proc. Lond. Math. Soc. (3) **6** (1956), no. 3, 366–416.
- [19] S. A. Huq, *Commutator, nilpotency and solvability in categories*, Q. J. Math. **19** (1968), no. 2, 363–389.
- [20] G. Janelidze, L. Márki, and W. Tholen, *Semi-abelian categories*, J. Pure Appl. Algebra **168** (2002), no. 2-3, 367–386.
- [21] P. T. Johnstone, *Sketches of an elephant: A topos theory compendium*, Oxford Logic Guides, vol. 43, 44, Oxford Univ. Press, 2002, Volumes 1 and 2.
- [22] R. Lavendhomme and J. R. Roisin, *Cohomologie non abélienne de structures algébriques*, J. Algebra **67** (1980), 385–414.
- [23] J.-L. Loday, *Une version non commutative des algèbres de Lie: les algèbres de Leibniz*, Enseign. Math. (2) **39** (1993), no. 3-4, 269–293.
- [24] S. Mac Lane, *Categories for the working mathematician*, second ed., Grad. Texts in Math., vol. 5, Springer, 1998.
- [25] N. Martins-Ferreira and T. Van der Linden, *A note on the "Smith is Huq" condition*, Appl. Categ. Structures **20** (2012), no. 2, 175–187.
- [26] G. Mason and G. Yamskulna, *Leibniz algebras and Lie algebras*, SIGMA Symmetry Integrability Geom. Methods Appl. **9** (2013), Paper 063, 10.
- [27] A. Montoli, *Action accessibility for categories of interest*, Theory Appl. Categ. **23** (2010), no. 1, 7–21.
- [28] G. Orzech, *Obstruction theory in algebraic categories I and II*, J. Pure Appl. Algebra **2** (1972), 287–314 and 315–340.
- [29] K. A. Zhevlakov, A. M. Slin'ko, I. P. Shestakov, and A. I. Shirshov, *Rings that are nearly associative*, Academic Press, 1982.

(Xabier García-Martínez) DEPARTMENT OF ALGEBRA, UNIVERSITY OF SANTIAGO DE COMPOSTELA, 15782 SANTIAGO DE COMPOSTELA, SPAIN.

E-mail address: `xabier.garcia@usc.es`

(Tim Van der Linden) INSTITUT DE RECHERCHE EN MATHÉMATIQUE ET PHYSIQUE, UNIVERSITÉ CATHOLIQUE DE LOUVAIN, CHEMIN DU CYCLOTRON 2 BTE L7.01.02, B-1348 LOUVAIN-LA-NEUVE, BELGIUM

E-mail address: `tim.vanderlinden@uclouvain.be`